

phenomenon and may not lead initially to enlargement of the liver.

Given the worsening nature of the patient's ascites after vitamin A intake was discontinued, a way was sought to maximize the removal of the vitamin A by increasing the RBP levels. Nutritional supplementation was given to provide an adequate protein intake. A trial of prednisone was used to increase the mobilization of hepatic vitamin A by increasing the release of RBP from the liver.^{1,12}

In summary, this case report describes an unusual presentation of chronic vitamin A intoxication. It suggests that children with malnutrition may be at a greater risk for intoxication because of impaired RBP production and can present with normal vitamin A levels. Determining the retinol-binding protein level is vital in the diagnosis and helpful in deciding the management. Last, although most patients' symptoms resolve simply by withdrawing the vitamin A supplementation and providing good nutrition, a trial of steroids may be considered in patients with persistent clinical symptoms.

REFERENCES

1. Smith JE, Goodman DS: Retinol-binding protein and the regulation of vitamin A transport. *Fed Proc* 1979; 38:2504-2509
2. Oliver TK Jr: Chronic vitamin A intoxication. *Am J Dis Child* 1958; 95:57-68
3. Muentner MD, Perry HO, Ludwig J: Chronic vitamin A intoxication in adults—Hepatic, neurologic and dermatologic complications. *Am J Med* 1971; 50:129-136
4. Farris WA, Erdman JW Jr: Protracted hypervitaminosis A following long-term, low-level intake. *JAMA* 1982; 247:1317-1318
5. Lippe B, Hensen L, Mendoza G, et al: Chronic vitamin A intoxication—A multisystem disease that could reach epidemic proportions. *Am J Dis Child* 1981; 135:634-636
6. Rosenberg HK, Berezin S, Heyman S, et al: Pleural effusion and ascites. *Clin Pediatr (Phila)* 1982; 21:435-440
7. Nosedá A, Alder M, Ketelbant P, et al: Massive vitamin A intoxication with ascites and pleural effusion. *J Clin Gastroenterol* 1985; 7:344-349
8. Silverman SH, Lecks HI: Protein-calorie deficiency and vitamin indiscretion in an atopic child who developed hypervitaminosis A. *Clin Pediatr (Phila)* 1982; 21:172-174
9. Guarascio P, Portmann B, Visco G, et al: Liver damage with reversible portal hypertension from vitamin A intoxication: Demonstration of Ito cells. *J Clin Pathol* 1983; 36:769-771
10. Russell RM, Boyer JL, Bagheri SA, et al: Hepatic injury from chronic hypervitaminosis A resulting in portal hypertension and ascites. *N Engl J Med* 1974; 291:435-440
11. Smith FR, Goodman DS: Vitamin A transport in human vitamin A toxicity. *N Engl J Med* 1976; 294:805-808
12. Atukorala TMS, Basu TK, Dickerson JWT: Effect of corticosterone on the plasma and tissue concentrations of vitamin A in rats. *Ann Nutr Metab* 1981; 25:234-238

Dysgonic Fermenter-2 Infections

RAMON E. PEREZ, MD
Anaheim, California

DOG BITES have been known to cause dangerous sepsis. The most commonly known bacterium is *Pasteurella multocida*. A search of the English language literature was made using MEDLINE 1977-1986 files. Since 1976 there have been multiple reports of dysgonic fermenter (DF-2) infections, many of them following dog bites or casual exposure to dogs. Dysgonic (Greek *dys*, "difficult," plus *gonikos*, "seed, offspring") fermenters are fastidious gram-negative rods easily missed by

ABBREVIATIONS USED IN TEXT

DF-2 = dysgonic fermenter
HIA = heart infusion agar

routine culture media. They are found in the oropharynx of some dogs. Patients with the worst prognosis tend to have splenectomies, alcoholism, or chronic obstructive pulmonary disease.

There have been 42 previously documented cases of DF-2 infections (one of them presumptive). The clinical spectrum has ranged from self-limited disease to fatal sepsis. The actual incidence is predictably higher than reported because of the difficulties with culture. DF-2 needs enriched media for growth and fermentation. Previous strains have reportedly been susceptible to penicillin. This is the first case description of penicillin resistance with DF-2.

Report of a Case

The patient, a 65-year-old man, had suffered from chronic obstructive pulmonary disease, nephrolithiasis, and hypertension. He was bitten on the left third finger by a dog on March 9, 1986. The following day he described shaking chills and on March 11 had temperatures up to 40°C (104°F). He was admitted to hospital the next day with a blood pressure of 85/70 mm of mercury, confusion, and fecal incontinence. The skin was acrocyanotic with ecchymotic, purpuric, and petechial lesions involving the extremities, face, and periumbilical area. Gangrene developed on the bitten finger without evidence of osteomyelitis. There were no mucosal or retinal lesions. The lungs were clear, and the heart had a II/VI systolic murmur at the apex and aortic valve area. The abdomen was soft without tenderness or organomegaly. There was no costovertebral angle tenderness despite complaints of back pain.

Initially the prothrombin time was 22 seconds, partial thromboplastin time 100 seconds, platelet count 14×10^9 per liter (14,000 per μ l), hematocrit 0.39 (38.7%), and hemoglobin 8.4 mmol per liter (13.5 grams per dl). The leukocyte count was 6.6×10^9 per liter (6,600 per μ l), with 0.54 (54%) neutrophils, 0.32 (32%) bands, 0.07 (7%) lymphocytes, and 0.06 (6%) metamyelocytes. Serum electrolyte values were normal except for a bicarbonate of 12 mmol per liter (12 mEq per liter). The blood urea nitrogen level was 22.1 mmol per liter (62 mg per dl) and the serum creatinine level 475 μ mol per liter (4.2 mg per dl).

The patient was treated with a regimen of penicillin G benzathine, 12 million units per day, and ceftizoxime sodium, 6 grams per day, given intravenously. He became more alert, and his hemodynamic indicators improved after fluid replacement and correction of the metabolic acidosis.

On March 15, he had a respiratory arrest requiring assisted ventilation. His condition continued to deteriorate, and he died on March 18, nine days after the dog bite. The main autopsy findings included a hemorrhagic spleen, congestion of the entire mucosa of the small and large intestines, and mucosal edema and hemorrhage of the major bronchi. The thyroid, adrenal, and pituitary glands were edematous.

All the blood cultures from admission became positive in 48 hours (March 14), with a fastidious gram-negative rod by the Bactec 6A automated system (Johnston Laboratories, Townson, Md). A preliminary identification of DF-2 was made on March 17, 1986, and was later corroborated by the State of California Microbial Diseases Laboratory. Its growth

(Perez RE: Dysgonic fermenter-2 infections. *West J Med* 1988 Jan; 148:90-92)

From the Division of Infectious Diseases, Department of Medicine, University of California, Irvine, California College of Medicine.

Reprint requests to Ramon E. Perez, MD, Infectious Disease Department, University of California, Irvine, 50 S Anaheim Blvd, #244, Anaheim, CA 92805.

requirements and biochemical pattern (Table 1) are similar to the previously reported DF-2. The organism does not grow on MacConkey agar or unsupplemented broth. Growth occurred in enriched chocolate agar (gonococcal agar, hemoglobin, reagents for *Neisseria* and *Hemophilus influenzae*) and blood agar (tryptic soy agar plus 5% sheep erythrocytes), both supplied by Remel Laboratories, Lenexa, Kansas. Repeat cultures after antibiotic therapy was begun remained negative.

The antibiotic susceptibility pattern is different from that in previously reported cases of DF-2 infection. Because of poor growth on routine media, the broth-disk elution method for anaerobic bacteria described by Wilkins and Thiel¹ was used. This patient's bacteria were susceptible to chloramphenicol (12 µg per ml), ceftizoxime (18 µg per ml), and metronidazole (16 µg per ml), but not to penicillin (2 units per ml), clindamycin (1.6 µg per ml), or erythromycin (3 µg per ml).

Discussion

The first report of a case of DF-2 infection was published in 1976 by Bobo and Newton.² The following year 17 cases, including that of Bobo and Newton, were reviewed by Butler and co-workers.³ An additional 25 cases have been reported in the English literature.⁴⁻²⁶

Most reviews of animal bites fail to recognize DF-2 as a pathogen. In a previous evaluation of canine oral flora, DF-2 was not detected.²⁷ In two reported cases DF-2 was recovered from the dogs' gingivae.^{8,26}

The clinical presentations of documented cases include self-limited infections without the need of antibiotics,⁹ cellulitis, gangrene, blepharitis,²⁵ arthritis, endocarditis¹⁵ without fevers,¹⁰ meningitis,^{2,12,26} brain abscess,¹⁶ disseminated intravascular coagulation, the Waterhouse-Friderichsen syndrome,¹¹ myocardial infarction,²³ and cardiopulmonary arrest.

Of the 43 cases, 24 (56%) were directly related to dog bites and others were associated with dogs. An underlying disease was present in 33 patients (77%). Splenectomy was noted in 17 patients (40%) and alcoholism in 10 (23%).

The overall mortality is 12/43 (28%). Nine of the deaths (9/12, or 75%) occurred in patients with some underlying disease. Conversely, 24 of 33 patients (73%) with underlying illness survived. Three of ten patients (30%) without underlying illness died. In four of ten patients (40%) who died, smears of their peripheral blood or buffy coats showed bacteria. There is no apparent element that correlates with the prognosis. It remains to be proved if the fatal cases are due to penicillin resistance or to a more virulent strain of DF-2.

The optimal culture medium for DF-2 has not been determined. The use of enriched chocolate and blood agar facilitated growth of the strain reported here. Fermentation reactions were possible by adding rabbit serum to the broth (Table 1). Different blood culture media have been used to recover DF-2. These include Bactec 6A, 6B aerobic bottles, 7C anaerobic bottles, trypticase-soy with sodium polyanethol sulfonate, supplemented peptone, and glucose broths. Subculture media have included heart infusion agar (HIA) with 5% rabbit blood in CO₂, trypticase-soy agar with 5% sheep blood, and HIA with 5% sheep blood.^{28,29}

Penicillin has been described as the drug of choice, but, as with this patient, one must be aware of resistant strains that until now had not been described. The routine methods for antibiotic susceptibility tests—Kirby-Bauer, minimum inhib-

TABLE 1.—Characteristics of Bacteria in Patient With Dysgonic Fermenter-2 Infection

Morphology	Long, thin, spindled gram-negative rod
Catalase	+
Oxidase	+
Growth	
MacConkey agar	—
SS agar	—
Blood and chocolate agar	
25°C (days)	+ (2-4)
35°C (days)	+ (2)
42°C	—
Aerobically	+
Carbon dioxide 5%	+
Fermentation	
Phenol red broth + rabbit serum	
Glucose (days)	A (2)
Lactose (days)	A (3-5)
Maltose (days)	A (3-5)
Triple sugar iron	
Slant/aerobic	Aw
Butt/anaerobic	—
Hydrogen sulfide	—
Lead acetate	+

A = acid; Aw = acid, weak reaction

itory concentration with Mueller-Hinton broth, and so forth—are standardized for rapidly growing aerobic or facultative organisms. Slow-growing bacteria like DF-2 require modified techniques that allow adequate growth of the organism to detect any effect by the antibiotic being tested. The original strains of DF-2 described by Butler and associates³ were tested by an agar dilution method using a special medium (Schaedler's). In most of the subsequent reports on DF-2, the authors failed to do or to report their susceptibility tests, sometimes only quoting the previous results of Butler and colleagues. Many community hospitals are not properly staffed or familiar with the technique of diluting agar for fastidious bacteria.

Broth-disk elution tests, such as Wilkins-Thiel's, are easier to do, are reproducible, and are more acceptable to laboratories with limited resources. Antibiotic disk(s) are diluted with broth in individual tubes. For penicillin, a 10-unit disk is diluted with 5 ml of medium, giving a penicillin concentration of 2 units per ml. One drop of bacterial inoculum is added to each tube containing antibiotic and to a control tube without antibiotic. The tubes are incubated and their turbidity is compared with the control tube. Susceptibility can be defined as the absence of turbidity. Reports are given as either susceptible or resistant to the given concentration of antibiotic tested.

Using penicillin alone might not be adequate therapy for sepsis following a dog bite. Administering a second antibiotic active against *Staphylococcus* and other gram-negative bacteria is appropriate while awaiting susceptibility studies. A reasonable regimen to treat a suspected DF-2 infection would be penicillin plus a third-generation cephalosporin like cefotaxime or ceftizoxime while awaiting bacteriologic reports. It would be difficult to justify the use of a first-generation cephalosporin as the main therapy for DF-2, considering its clinical failure with *P. multocida*, the lack of central nervous system penetration, and the possible diagnoses listed below.

When a patient presents with petechiae, purpura, and ec-

chymosis, the differential diagnosis includes meningococemia, Rocky Mountain spotted fever, and sepsis in asplenic and immune-compromised patients. The history of a dog bite should suggest the possibility of *P. multocida* and DF-2 infections. Penicillin appears to be the drug of choice for DF-2 infections, but susceptibility tests should be done on all clinical organisms for the possibility of a resistant strain. The best alternative for patients who are either allergic to penicillin or who have penicillin-resistant infections is not known.

REFERENCES

1. Wilkins TD, Thiel T: Modified broth-disk method for testing the antibiotic susceptibility of anaerobic bacteria. *Antimicrob Agents Chemother* 1973; 3:350-356
2. Bobo RA, Newton EJ: A previously undescribed gram-negative bacillus causing septicemia and meningitis. *Am J Clin Pathol* 1976; 65:564-569
3. Butler T, Weaver RE, Ramani TKV, et al: Unidentified gram-negative rod infection—A new disease of man. *Ann Intern Med* 1977; 86:1-5
4. Schlossberg D: Septicemia caused by DF-2. *J Clin Microbiol* 1979; 9:297-298
5. Boyce F, Price RF: Splenectomized patient with fatal DF-2 septicemia. *Clin Microb Newsletter* 1980; 2:4
6. Findling JW, Pohlmann GP, Rose HD: Fulminant gram-negative bacillemia (DF-2) following a dog bite in an asplenic woman. *Am J Med* 1980; 68:154-156
7. Hinrichs JH, Dunkelberg WE: DF-2 septicemia after splenectomy: Epidemiology and immunologic response. *South Med J* 1980; 73:1638-1640
8. Martone WJ, Zuehl RW, Minson GE, et al: Postsplenectomy sepsis with DF-2: Report of a case with isolation of the organism from the patient's dog. *Ann Intern Med* 1980; 93:457-458
9. Schoen RT, Wohlgeleit D, Barden GE, et al: Infection with CDC group DF-2 gram-negative rod: Report of two cases. *Arch Intern Med* 1980; 140:657-658
10. Shankar PS, Scott JH, Anderson CL: Atypical endocarditis due to gram-negative bacillus transmitted by dog bite. *South Med J* 1980; 73:1640-1641
11. Chaudhuri AK, Hartley RB, Maddocks AC: Waterhouse-Friderichsen syndrome caused by a DF-2 bacterium in a splenectomized patient. *J Clin Pathol* 1981; 34:172-173
12. Ofori-Adjei D, Blackledge P, O'Neill P: Meningitis caused by dysgonic fermenter type 2 (DF-2) organism in a previously healthy adult. *Br Med J [Clin Res]* 1982; 285:263-264
13. Smith C, Rubin SJ: DF-2 septicemia. *Clin Microb Newsletter* 1983; 5:90-91
14. Howard AJ, Hughes M, Parry H, et al: Dog bites and dysgonic fermenting organisms (Letter). *Lancet* 1983; 2:1022-1023
15. Worthington M, Gleason T, Pandian NG, et al: Tricuspid valve myxoma infected with dysgonic fermenter-2. *South Med J* 1984; 77:241-242
16. Archer SL: Dysgonic fermenter 2 infection resulting in chronic glomerulonephritis. *Can Med Assoc J* 1985; 132:657-660
17. Fibbe W, Lighart G, van den Broek P, et al: Septicemia with a dysgonic fermenter-2 (DF-2) bacterium in a compromised host. *Infection* 1985; 13:286-287
18. Kalb R, Kaplan MH, Tenenbaum MJ, et al: Cutaneous infection at dog bite wounds associated with fulminant DF-2 septicemia. *Am J Med* 1985; 78:687-690
19. Scully RE, Mark EJ, McNeely BU: Case records of the Massachusetts General Hospital, Case 29-1986. *N Engl J Med* 1986; 315:241-249
20. Pers C, Kristiansen JE, Scheibel JH, et al: Fatal septicemia caused by DF-2 in a previously healthy man. *Scand J Infect Dis* 1986; 18:265-267
21. Waters MJ, Stern L, Andrew JH: Dysgonic fermenter-2 septicemia. *Pathology* 1986; 18:162-164
22. Holmes RL, Kozinn WP: DF-2 septicemia following whirlpool spa immersion. *J Clin Microbiol* 1986; 23:627-628
23. Newton NL, Sharma B: Acute myocardial infarction associated with DF-2 bacteremia after a dog bite. *Am J Med Sci* 1986; 291:352-354
24. Tison DL, Latimer JM: Lysis-centrifugation-direct plating technique for isolation of group DF-2 from the blood of a dog bite victim [Letter]. *J Infect Dis* 1986; 153:1001-1002
25. Glasser DB: Angular blepharitis caused by gram-negative bacillus DF-2. *Am J Ophthalmol* 1986; 102:119-120
26. Chan PC, Fonseca K: Septicaemia and meningitis caused by dysgonic fermenter-2 (DF-2). *J Clin Pathol* 1986; 39:1021-1024
27. Bailie WE, Stowe EC, Schmitt AM: Aerobic bacterial flora of oral and nasal fluids of canines with reference to bacteria associated with bites. *J Clin Microbiol* 1978; 7:223-231
28. Weaver RE, Hollis DG, Bottone EJ: Gram-negative fermentative bacteria and *Francisella tularensis*, in Lennette EH, Balows A, Hausler WJ, Jr, et al (Eds): *Manual of Clinical Microbiology*, 4th Ed. Washington, DC, American Society for Microbiology, 1985, pp 325-326
29. Weaver RE: DF-2. *Microbiology* 1985; 28:1-5